IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of: Takashi KADOWAKI et al.

Serial No.:

10/591,490

Group No.:

Filed:

09/01/2006

Examiner:

For:

Agents For Regulating Adiponectin Receptor Expression

INFORMATION DISCLOSURE STATEMENT

MS PCT

Commissioner for Patents P.O. Box 1450 Alexandria, VA 22313-1450

CERTIFICATE OF MAILING UNDER 37 CFR § 1.8(a)

I hereby certify that this correspondence is being deposited with the United States Postal Service as first class mail in an envelope addressed to the: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450, on May 8, 2007.

Dear Sir or Madam:

The citations listed below, copies attached, may be material to the examination of the above-identified application, and are therefore submitted in compliance with the duty of disclosure defined in 37 C.F.R. § 1.56 and § 1.97. The Examiner is requested to make these citations of official record in this application:

- Publication No. US 2004/0241802 A1 of Kadowaki et al., "Adiponectin receptor and gene encoding same," (2004) as English translation of related PCT application;
- Publication No. WO 2004/061108 A1 of Kadowaki et al., "Adiponectin receptor and gene encoding same," (2004) in Japanese accompanied by English translation of related national entry application;
- Berg et al., "The adipocyte-secreted protein Acrp30 enhances hepatic insulin action," Nature Medicine, 7:947-953 (2001);
- Biggs III et al., "Protein kinase B/Akt-mediated phosphorylation promotes nuclear exclusion of the winged helix transcription factor FKHR1," Proc Natl Acad Sci USA, 96:7421-7426 (1999);

- Brunet *et al.*, "Akt promotes cell survival by phosphorylating and inhibiting a forkhead transcription factor," *Cell.* 96:857-868 (1999);
- Friedman et al., "Phosphoenolpyruvate carboxykinase (GTP) gene transcription and hyperglycemia are regulated by glucocorticoids in genetically obese db/db transgenic mice," J Biol Chem, 272:31475-31481 (1997);
- Fruebis et al., "Proteolytic cleavage product of 30-kDa adipocyte complement-related protein increases fatty acid oxidation in muscle and causes weight loss in mice," Proc Natl Acad Sci USA, 98:2005-2010 (2001);
- Guo *et al.*, "Phosphorylation of serine 256 by protein kinase B disrupts transactivation by FKHR and mediates effects of insulin on insulin-like growth factor-binding protein-1 promoter activity through a conserved insulin response sequence," *J Biol Chem*, 274:17184-17192 (1999);
- Herzig *et al.*, "CREB regulates hepatic gluconeogenesis through the coactivator PGC-1," *Nature*, 413:179-183 (2001);
- Hu et al., "AdipoQ is a novel adipose-specific gene dysregulated in obesity," *J Biol Chem*, 271:10697-10703 (1996);
- Kadowaki, "Insights into insulin resistance and type 2 diabetes from knockout mouse models," *J Clin Invest*, 106:459-465 (2000);
- Kubota *et al.*, "Disruption of adiponectin causes insulin resistance and neointimal formation," *J Biol Chem*, 277:25863-25866 (2002);
- Levine et al., "Toxicologic evaluation of streptozotocin (NSC 85998) in mice, dogs and monkeys," *Drug Chem Toxicol*, 3:201-212 (1980);
- Maeda et al., "cDNA cloning and expression of a novel adipose specific collagen-like factor, apM1 (Adipose Most Abundant Gene Transcript 1), Biochem Biophys Res Commun, 221:286-289 (1996);
- Maeda et al., "Diet-induced insulin resistance in mice lacking adiponectin/ACRP30,"
 Nature Medicine, 8:731-737 (2002);
- Nakae et al., "Insulin stimulates phosphorylation of the forkhead transcription factor
 FKHR on serine 253 through a wortmannin-sensitive pathway," J Biol Chem, 274:1598215985 (1999);

- Nakae et al., "The forkhead transcription factor Foxo1 (Fkhr) confers insulin sensitivity onto glucose-6-phosphatase expression," J Clin Invest, 108:1359-1367 (2001);
- Nakano et al., "Isolation and characterization of GBP28, a novel gelatin-binding protein purified from human plasma," J Biochem, 120:803-812 (1996);
- Ouchi et al., "Adipocyte-derived plasma protein, adiponectin, suppresses lipid accumulation and class A scavenger receptor expression in human monocyte-derived macrophages," Circulation, 103:1057-1063 (2001);
- Rakieten *et al.*, "Studies on the diabetogenic action of streptozotocin (NSC-37917)," *Cancer Chemother Rep*, 29:91-98 (1963);
- Scheer *et al.*, "Constitutively active mutants of the α1B-adrenergic receptor: role of highly conserved polar amino acids in receptor action," *EMBO J*, 15:3566-3578 (1996);
- Scherer et al., "A novel serum protein similar to C1q, produced exclusively in adipocytes," J Biol Chem, 270:26746-26749 (1995);
- Seglen, "Preparation of isolated rat liver cells," *Methods Cell Biol*, 13:29-83 (1976);
- Shepherd *et al.*, "Phosphoinositide 3-kinase: the key switch mechanism in insulin signaling," *Biochem J*, 333:471-490 (1998);
- Takaishi *et al.*, "Regulation of nuclear translocation of Forkhead transcription factor AFX by protein kinase B," *Proc Natl Acad Sci USA*, 96:11836-11841 (1999);
- Tomas *et al.*, "Enhanced muscle fat oxidation and glucose transport by ACRP30 globular domain: Acetyl-CoA carboxylase inhibition and AMP-activated protein kinase activation," *Proc Natl Acad Sci USA*, 99:16309-16313 (2002);
- Tsuchida *et al.*, "Insulin/Foxo1 pathway regulates expression levels of adiponectin receptors and adiponectin sensitivity," *J Biol Chem*, 279:30817-30822 (2004);
- Virkamaki *et al.*, "Protein-protein interaction in insulin signaling and the molecular mechanisms of insulin resistance," *J Clin Invest*, 108:1001-1013 (2001);
- Wess, "G-protein-coupled receptors: Molecular mechanisms involved in receptor activation and selectivity of G-protein recognition," FASEB J, 11:346-354 (1997);
- Yamauchi et al., "The fat-derived hormone adiponectin reverses insulin resistance associated with both lipoatrophy and obesity," *Nature Medicine*, 7:941-946 (2001);

- Yamauchi et al., "Inhibition of RXR and PPARγ ameliorates diet-induced obesity and type 2 diabetes," J Clin Invest, 108:1001-1013 (2001);
- Yamauchi et al., "Adiponectin stimulates glucose utilization and fatty-acid oxidation by activating AMP-activated protein kinase," *Nature Medicine*, 8:1288-1295 (2002);
- Yamauchi et al., "Globular adiponectin protected ob/ob mice from diabetes and ApoE-deficient mice from atherosclerosis," J Biol Chem, 278:2461-2468 (2003);
- Yamauchi et al., "Cloning of adiponectin receptors that mediate antidiabetic metabolic effects," Nature, 324:762-769 (2003); and
- Yokomizo *et al.*, "A G-protein-coupled receptor for leukotriene B₄ that mediates chemotaxis," *Nature*, 387:620-624 (1997).

This Information Disclosure Statement under 37 C.F.R. § 1.56 and § 1.97 is not to be construed as a representation that a search has been made, that additional information material to the examination of this application does not exist, or that any one or more of these citations constitutes prior art.

Dated: May 8, 2007

Christine A. Lekutis Registration No. 51,934

MEDLEN & CARROLL, LLP 101 Howard Street, Suite 350 San Francisco, California 94105 415/904-6500 FORM PTO-1449 (Modified)

U.S. Department of Commerce Patent and Trademark Office

| Attorney Docket No.: SHIMIZU-13116 | П | Serial No · | 10/591 490 |
|---------------------------------------|---|-------------|------------|
| Attorney Docket No.: Smilling 0-13110 | ┸ | Senai No.: | 10/391,490 |

Applicant: Takashi Kadowaki

| INFORMATION DISCLOSURE STATEMENT BY APPLICANT | | | Applicant. Takashi Kadi | UWAKI | | | | | | |
|---|---------------|--|-----------------------------|---------------------------|-------------------------------|--------------------|-----------------|-------------|--------------------|--|
| (Use Several Sheets If Necessary) (37 CFR § 1.98(b)) Filing or 371(c) Date: (| | | | /01/2006 Group Art Unit: | | | | | | |
| | | | | U.S. PATENT DOC | UMENTS | | | | | |
| Examiner Initials | Cite No. | Document / Patent Number | Publication / Issue Date | Applic | Applicant / Patentee | | Subclass | Filing Date | | |
| | 1 | 2004/0241802 | | Kadowaki et al. | | | | | | |
| <u>.</u> | | | | | | | | | | |
| . | | F | OREIGN PATENTS | OR PUBLISHED FOR | REIGN PATENT APPLICA | TIONS | | | | |
| • | | Document Number | Publication Date | Country | Country / Patent Office | | Subclass | | Translation Yes No | |
| | 2 | WO 2004/061108 | | PCT | | | | 165 | INO | |
| | | | | 101 | | | | | | |
| | | OTHER | DOCUMENTS (Incl | uding Author, Title, D | ate, Relevant Pages, Place of | of Publication) | • | | | |
| | 3 | Berg et al., "The adi | pocyte-secreted protei | in Acrp30 enhances he | patic insulin action," Natur | e Medicine, 7:94 | 7-953 (2001) | | | |
| | 4 | Biggs III et al., "Protein kinase B/Akt-mediated phosphorylation promotes nuclear exclusion of the winged helix transcription factor FKHR1," Proc Natl Acad Sci USA, 96:7421-7426 (1999) | | | | | | | | |
| | 5 | Brunet et al., "Akt p | romotes cell survival | by phosphorylating an | d inhibiting a forkhead trans | scription factor," | Cell, 96:857-86 | 8 (1999) | | |
| | 6 | Brunet et al., "Akt promotes cell survival by phosphorylating and inhibiting a forkhead transcription factor," Cell, 96:857-868 (1999) Friedman et al., "Phosphoenolpyruvate carboxykinase (GTP) gene transcription and hyperglycemia are regulated by glucocorticoids in genetically obese db/db transgenic mice," J Biol Chem, 272:31475-31481 (1997) | | | | | | | | |
| | 7 | Fruebis et al., "Proteolytic cleavage product of 30-kDa adipocyte complement-related protein increases fatty acid oxidation in muscle and causes weight loss in mice," Proc Natl Acad Sci USA, 98:2005-2010 (2001) | | | | | | | | |
| | 8 | Guo et al., "Phosphorylation of serine 256 by protein kinase B disrupts transactivation by FKHR and mediates effects of insulin on insulin-like growth factor-binding protein-1 promoter activity through a conserved insulin response sequence," J Biol Chem, 274:17184-17192 (1999) | | | | | | | | |
| | 9 | Herzig et al. "CREB regulates hepatic gluconeogenesis through the coactivator PGC-1," Nature, 413:179-183 (2001) | | | | | | | | |
| | 10 | Hu et al., "AdipoQ is a novel adipose-specific gene dysregulated in obesity," J Biol Chem, 271:10697-10703 (1996) | | | | | | | | |
| | 11 | Kadowaki, "Insights into insulin resistance and type 2 diabetes from knockout mouse models," J Clin Invest, 106:459-465 (2000) | | | | | | | | |
| | 12 | Kubota et al., "Disru | ption of adiponectin c | causes insulin resistance | e and neointimal formation | ," J Biol Chem, i | 277:25863-2586 | 6 (2002) | | |
| | 13 | Kubota et al., "Disruption of adiponectin causes insulin resistance and neointimal formation," J Biol Chem, 277:25863-25866 (2002) Levine et al., "Toxicologic evaluation of streptozotocin (NSC 85998) in mice, dogs and monkeys," Drug Chem Toxicol, 3:201-212 (1980) | | | | | | | | |
| | 14 | Maeda et al., "cDNA cloning and expression of a novel adipose specific collagen-like factor, apM1 (Adipose Most Abundant Gene Transcript 1), Biochem Biophys Res Commun, 221:286-289 (1996) | | | | | | | | |
| | 15 | Maeda et al., "Diet-induced insulin resistance in mice lacking adiponectin/ACRP30," Nature Medicine, 8:731-737 (2002) | | | | | | | | |
| | 16 | Nakae et al., "Insulin stimulates phosphorylation of the forkhead transcription factor FKHR on serine 253 through a wortmannin-sensitive pathway," J Biol Chem, 274:15982-15985 (1999) | | | | | | | | |
| | 17 | Nakae et al., "The forkhead transcription factor Foxol (Fkhr) confers insulin sensitivity onto glucose-6-phosphatase expression," J Clin Invest, 108:1359-1367 (2001) | | | | | | | | |
| | 18 | Nakano et al., "Isolation and characterization of GBP28, a novel gelatin-binding protein purified from human plasma," J Biochem, 120:803-812 (1996) | | | | | | | | |
| | 19 | Ouchi et al., "Adipocyte-derived plasma protein, adiponectin, suppresses lipid accumulation and class A scavenger receptor expression in human monocyte-derived macrophages," Circulation, 103:1057-1063 (2001) | | | | | | | | |
| | 20 | Rakieten et al., "Studies on the diabetogenic action of streptozotocin (NSC-37917)," Cancer Chemother Rep, 29:91-98 (1963) | | | | | | | | |
| | 21 | Scheer et al., "Constitutively active mutants of the α1B-adrenergic receptor: role of highly conserved polar amino acids in receptor action," EMBO J, 15:3566-3578 (1996) | | | | | | | | |
| | 22 | Scherer et al., "A novel serum protein similar to C1q, produced exclusively in adipocytes," J Biol Chem, 270:26746-26749 (1995) | | | | | | | | |
| | 23 | Seglen, "Preparation of isolated rat liver cells," Methods Cell Biol, 13:29-83 (1976) | | | | | | | | |
| | 24 | Shepherd et al., "Phosphoinositide 3-kinase: the key switch mechanism in insulin signaling," Biochem J, 333:471-490 (1998) | | | | | | | | |
| | 25 | Takaishi et al., "Regulation of nuclear translocation of Forkhead transcription factor AFX by protein kinase B," Proc Natl Acad Sci USA, 96:11836-11841 (1999) | | | | | | | | |
| Examiner: | | | | | Date Considered: | | | | | |
| EVAMINED. | Initial citat | ion considered Draw | line through citation is | f not in conformers = | nd not considered. Include | aanu af thia fa | n udeh maue | | | |

EXAMINER: Initial citation considered. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

| | | | Sheet 2 of | | | |
|--|---|---|------------------------|--|--|--|
| FORM PTO-1449 (Modified) | U.S. Department of Commerce Patent and Trademark Office | Attorney Docket No.: SHIMIZU-13116 | Serial No.: 10/591,490 | | | |
| 1 ` ′ | | Applicant: Takashi Kadowaki | | | | |
| INFORMATION DISCLOSURE STATEMENT BY APPLICANT (Use Several Sheets If Necessary) (37 CFR § 1.98(b)) | | Filing or 371(c) Date: 09/01/2006 | Group Art Unit: | | | |
| | OTHER DOCUMENTS (Including Author, Title, D | Date, Relevant Pages, Place of Publication) | | | | |
| 26 | Tomas et al., "Enhanced muscle fat oxidation and glucose transport by ACRP30 globular domain: Acetyl-CoA carboxylase inhibition and AMP-activated protein kinase activation," Proc Natl Acad Sci USA, 99:16309-16313 (2002) | | | | | |
| 27 | Tsuchida et al., "Insulin/Foxo1 pathway regulates expression levels of adiponectin receptors and adiponectin sensitivity," J Biol Chem, 279:30817-30822 (2004) | | | | | |
| - 28 | Virkamaki et al., "Protein-protein interaction in insulin signaling and the molecular mechanisms of insulin resistance," J Clin Invest, 108:1001-1013 (2001) | | | | | |
| . 29 | Wess, "G-protein-coupled receptors: Molecular mechanisms involved in receptor activation and selectivity of G-protein recognition," FASEB J, 11:346-354 (1997) | | | | | |
| 30 | Yamauchi et al., "The fat-derived hormone adiponectin reverses insulin resistance associated with both lipoatrophy and obesity," Nature Medicine, 7:941-946 (2001) | | | | | |
| 31 | Yamauchi et al., "Inhibition of RXR and PPARy ameliorates diet-induced obesity and type 2 diabetes," J Clin Invest, 108:1001-1013 (2001) | | | | | |
| 32 | Yamauchi et al., "Adiponectin stimulates glucose utilization and fatty-acid oxidation by activating AMP-activated protein kinase," Nature Medicine, 8:1288-1295 (2002) | | | | | |
| 33 | Yamauchi et al., "Globular adiponectin protected ob/ob mice from diabetes and ApoE-deficient mice from atherosclerosis," J Biol Chem, 278:2461-2468 (2003) | | | | | |
| 34 | Yamauchi et al., "Cloning of adiponectin receptors that mediate antidiabetic metabolic effects," Nature, 324:762-769 (2003) | | | | | |
| 35 | Yokomizo et al., "A G-protein-coupled receptor for leukotriene B4 that mediates chemotaxis," Nature, 387:620-624 (1997) | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |

Examiner: Date Considered:

EXAMINER: Initial citation considered. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.